



## ON A SOLVABLE SYSTEM OF DIFFERENCE EQUATIONS OF HIGHER-ORDER WITH PERIOD TWO COEFFICIENTS

YASIN YAZLIK AND MERVE KARA

ABSTRACT. We show that the next difference equations system

$$x_{n+1} = \frac{a_n x_{n-k+1} y_{n-k}}{y_n - \alpha_n} + \beta_{n+1}, \quad y_{n+1} = \frac{b_n y_{n-k+1} x_{n-k}}{x_n - \beta_n} + \alpha_{n+1}, \quad n \in \mathbb{N}_0,$$

where  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ , the sequences  $(a_n)_{n \in \mathbb{N}_0}$ ,  $(b_n)_{n \in \mathbb{N}_0}$ ,  $(\alpha_n)_{n \in \mathbb{N}_0}$ ,  $(\beta_n)_{n \in \mathbb{N}_0}$  are two periodic and the initial conditions  $x_{-i}$ ,  $y_{-i}$   $i \in \{0, 1, \dots, k\}$ , are non-zero real numbers, can be solved. Also, we investigate the behavior of solutions of above mentioned system when  $a_0 = b_1$  and  $a_1 = b_0$ .

### 1. INTRODUCTION

Theory of difference equations have attracted attention of many authors in recent years(see, e.g., [1]-[40]). Most of the recent applications of this theory have appeared in many scientific areas such as biology, physics, engineering, economics. Particularly, rational difference equations and their systems of higher order have great importance in applications. It is very worthy to find systems which belong to solvable nonlinear difference equations systems and to solve nonlinear difference equations or systems in closed-form or explicit-form. The found formulas for the solutions of these types of equations or systems can be used easily for description of many features of the solutions of these equations or systems. For this reason, finding of a formula for solution of a nonlinear difference equation is worthy as well as interesting.

In an earlier paper, Elabbasy et al., in [3], considered, among other things, the next difference equation

$$x_{n+1} = \frac{x_n x_{n-1}}{x_n - 1} + 1, \quad n \in \mathbb{N}_0. \quad (1)$$

Received by the editors: January 11, 2018; Accepted: February 18, 2019.

2010 *Mathematics Subject Classification.* 39A10, 39A20, 39A23.

*Key words and phrases.* Periodic solution, solution in closed form system of difference equations.

Quite recently in [14], Haddad et al. considered the following system of difference equations

$$x_{n+1} = \frac{ax_n y_{n-1}}{y_n - \alpha} + \beta, \quad y_{n+1} = \frac{by_n x_{n-1}}{x_n - \beta} + \alpha, \quad n \in \mathbb{N}_0, \quad (2)$$

where the parameters  $a, b, \alpha, \beta$  and the initial conditions  $x_{-i}, y_{-i}, i = 0, 1$ , are non-zero real numbers, which is an extension of the equation in (1). By using appropriate substitutions on variables, authors reduced system (2) to the first order linear difference equations and investigated the existence and behavior of the solutions of system (2).

Our aim in this study to show that the next difference equations system

$$x_{n+1} = \frac{a_n x_{n-k+1} y_{n-k}}{y_n - \alpha_n} + \beta_{n+1}, \quad y_{n+1} = \frac{b_n y_{n-k+1} x_{n-k}}{x_n - \beta_n} + \alpha_{n+1}, \quad n \in \mathbb{N}_0, \quad (3)$$

where the sequences  $a_n, b_n, \alpha_n, \beta_n$  are two periodic and the initial conditions  $x_{-i}, y_{-i}, i \in \{0, 1, \dots, k\}$ , are non-zero real numbers, is solvable in closed form. Also, by using obtained formulas we give the behavior and periodicity of well-defined solutions of system (3) when  $a_0 = b_1$  and  $a_1 = b_0$ . Note that system (3) is a natural extension of both Eq. (1) and system (2),

**Lemma 1.** [27] *Let  $(a_n)_{n \in \mathbb{N}_0}$  and  $(b_n)_{n \in \mathbb{N}_0}$  be two sequences of real numbers and the sequences  $y_{km+i}, i = 0, k-1$ , be solutions of the equations*

$$y_{km+i} = a_{km+i} y_{k(m-1)+i} + b_{km+i}, \quad m \in \mathbb{N}_0. \quad (4)$$

*Then, for each fixed  $i \in \{0, 1, \dots, k-1\}$  and  $m \geq -1$ , equation (4) has the general solution*

$$y_{km+i} = y_{-k+i} \prod_{j=0}^m a_{kj+i} + \sum_{s=0}^m b_{ks+i} \prod_{j=s+1}^m a_{kj+i}.$$

*Further, if  $(a_n)_{n \in \mathbb{N}_0}$  and  $(b_n)_{n \in \mathbb{N}_0}$  are constant and  $i = 0, k-1$ , then*

$$y_{km+i} = \begin{cases} a^{m+1} y_{-k+i} + b \frac{1-a^{m+1}}{1-a}, & \text{if } a \neq 1, \\ y_{-k+i} + b(m+1), & \text{if } a = 1. \end{cases}$$

**Definition 2.** [12] *(Periodicity) A solution  $\{x_n\}_{n=-k}^{\infty}$  of equation  $x_{n+1} = f(x_n, x_{n-1}, \dots, x_{n-k})$ ,  $n \in \mathbb{N}_0$ , is called periodic with period  $p$  if there exists an integer  $p \geq 1$  such that  $x_{n+p} = x_n$  for all  $n \geq -k$ .*

In the sequel, as usual, we suppose that  $\prod_{j=i}^m A_j = 1$  and  $\sum_{j=i}^m A_j = 0$ , for all  $m < i$ .

## 2. SOLUTIONS OF SYSTEM (3)

Let  $(x_n, y_n)_{n \geq -k}$  be a solution of system (3). If one of the initial conditions  $x_{-j}, y_{-j}, j = 0, k$  is equal to zero, then the system (3) is not defined. For example, if  $x_{-k} = 0$ , then  $y_1 = \alpha_1$ , and so  $x_2$  can not be calculated. For the other  $j = 0, k-1$ ,

the case is same. Now assume that  $x_{-j} \neq 0 \neq y_{-j}$ ,  $j = \overline{0, k}$ . If one of the terms  $x_{n_0}$  and  $y_{n_0}$ , for  $n_0 \geq 1$ , is equal to zero, then from system (3) either  $x_{n_0+k} = \frac{a_{n_0+k-1}x_{n_0}y_{n_0-1}}{y_{n_0+k-1}-\alpha_{n_0+k-1}} + \beta_{n_0+k} = \beta_{n_0+k}$  and so, it follows that,  $y_{n_0+k+1}$  is not defined or  $y_{n_0+k} = \frac{b_{n_0+k-1}y_{n_0}x_{n_0-1}}{x_{n_0+k-1}-\beta_{n_0+k-1}} + \alpha_{n_0+k} = \alpha_{n_0+k}$  and so, it follows that,  $x_{n_0+k+1}$  is not defined. Thus, for well-defined solution of system (3), we may assume that none of the terms of  $x_n$  and  $y_n$  is not equal to zero, for every  $n \geq -k$ . By means of the change of variables

$$u_n = \frac{x_n - \beta_n}{x_{n-k}}, v_n = \frac{y_n - \alpha_n}{y_{n-k}}, n \in \mathbb{N}_0, \tag{5}$$

the system in (3) becomes

$$u_{n+1} = \frac{a_n}{v_n}, v_{n+1} = \frac{b_n}{u_n}, n \in \mathbb{N}_0. \tag{6}$$

From (6), we have two independent equations

$$u_{n+2} = \frac{a_{n+1}}{b_n}u_n, v_{n+2} = \frac{b_{n+1}}{a_n}v_n, n \in \mathbb{N}_0, \tag{7}$$

and so

$$u_{2m+i} = \left(\frac{a_{1-i}}{b_i}\right)^m u_i, v_{2m+i} = \left(\frac{b_{1-i}}{a_i}\right)^m v_i, i \in \{0, 1\}, m \in \mathbb{N}_0, \tag{8}$$

where the sequences  $(a_n)_{n \in \mathbb{N}_0}$  and  $(b_n)_{n \in \mathbb{N}_0}$  are two periodic. From (5) we have that

$$\begin{aligned} x_n &= u_n x_{n-k} + \beta_n = u_n u_{n-k} x_{n-2k} + u_n \beta_{n-k} + \beta_n, \\ y_n &= v_n y_{n-k} + \alpha_n = v_n v_{n-k} y_{n-2k} + v_n \alpha_{n-k} + \alpha_n, n \in \mathbb{N}_0. \end{aligned} \tag{9}$$

We consider three cases: (a)  $k = 1$ ; (b)  $k = 2t + 1$  ( $t = 1, 2, \dots$ ); and (c)  $k = (2t)$  ( $t = 1, 2, \dots$ ).

a) If  $k = 1$ , then, from equations in (9), we can write

$$x_n = u_n x_{n-1} + \beta_n, y_n = v_n y_{n-1} + \alpha_n, n \in \mathbb{N}_0. \tag{10}$$

from which it follows that

$$\begin{aligned} x_{2n} &= u_{2n} x_{2n-1} + \beta_{2n} = \left(\frac{a_1}{b_0}\right)^n u_0 x_{2n-1} + \beta_0, n \in \mathbb{N}_0, \\ x_{2n+1} &= u_{2n+1} x_{2n} + \beta_{2n+1} = \left(\frac{a_0}{b_1}\right)^n u_1 x_{2n} + \beta_1, n \in \mathbb{N}_0, \\ y_{2n} &= v_{2n} y_{2n-1} + \alpha_{2n} = \left(\frac{b_1}{a_0}\right)^n v_0 y_{2n-1} + \alpha_0, n \in \mathbb{N}_0, \\ y_{2n+1} &= v_{2n+1} y_{2n} + \alpha_{2n+1} = \left(\frac{b_0}{a_1}\right)^n v_1 y_{2n} + \alpha_1, n \in \mathbb{N}_0. \end{aligned}$$

which implies that

$$x_{2n+1} = \left(\frac{a_0}{b_1}\right)^n \left(\frac{a_1}{b_0}\right)^n u_0 u_1 x_{2n-1} + \left(\frac{a_0}{b_1}\right)^n u_1 \beta_0 + \beta_1, \quad n \in \mathbb{N}_0, \quad (11)$$

$$x_{2n+2} = \left(\frac{a_0}{b_1}\right)^n \left(\frac{a_1}{b_0}\right)^{n+1} u_0 u_1 x_{2n} + \left(\frac{a_1}{b_0}\right)^{n+1} u_0 \beta_1 + \beta_0, \quad n \in \mathbb{N}_0, \quad (12)$$

$$y_{2n+1} = \left(\frac{b_0}{a_1}\right)^n \left(\frac{b_1}{a_0}\right)^n v_0 v_1 y_{2n-1} + \left(\frac{b_0}{a_1}\right)^n v_1 \alpha_0 + \alpha_1, \quad n \in \mathbb{N}_0, \quad (13)$$

$$y_{2n+2} = \left(\frac{b_0}{a_1}\right)^n \left(\frac{b_1}{a_0}\right)^{n+1} v_0 v_1 y_{2n} + \left(\frac{b_1}{a_0}\right)^{n+1} v_0 \alpha_1 + \alpha_0, \quad n \in \mathbb{N}_0. \quad (14)$$

Hence, from Lemma 1 for  $k = 1$  and Eqs. (11)-(14), we can write the solution of system (3) as follows:

$$\begin{aligned} x_{2n-1} &= x_{-1} \prod_{t=0}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^t + \sum_{r=0}^{n-1} \beta_0 u_1 \left(\frac{a_0}{b_1}\right)^r \prod_{t=r+1}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^t \\ &\quad + \sum_{r=0}^{n-1} \beta_1 \prod_{t=r+1}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^t, \end{aligned}$$

$$\begin{aligned} x_{2n} &= x_0 \prod_{t=0}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^{t+1} + \sum_{r=0}^{n-1} \beta_0 \prod_{t=r+1}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^{t+1} \\ &\quad + \sum_{r=0}^{n-1} \beta_1 u_0 \left(\frac{a_1}{b_0}\right)^{r+1} \prod_{t=r+1}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^{t+1}, \end{aligned}$$

$$\begin{aligned} y_{2n-1} &= y_{-1} \prod_{t=0}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^t + \sum_{r=0}^{n-1} \alpha_0 v_1 \left(\frac{b_0}{a_1}\right)^r \prod_{t=r+1}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^t \\ &\quad + \sum_{r=0}^{n-1} \alpha_1 \prod_{t=r+1}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^t, \end{aligned}$$

$$\begin{aligned} y_{2n} &= y_0 \prod_{t=0}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^{t+1} + \sum_{r=0}^{n-1} \alpha_0 \prod_{t=r+1}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^{t+1} \\ &\quad + \sum_{r=0}^{n-1} \alpha_1 v_0 \left(\frac{b_1}{a_0}\right)^{r+1} \prod_{t=r+1}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^{t+1}, \end{aligned}$$

where  $u_0 = \frac{x_0 - \beta_0}{x_{-k}}$ ,  $u_1 = \frac{x_1 - \beta_1}{x_{1-k}}$ ,  $v_0 = \frac{y_0 - \alpha_0}{y_{-k}}$ ,  $v_1 = \frac{y_1 - \alpha_1}{y_{1-k}}$ ,  $d = u_0 u_1$  and  $v_0 v_1 = \frac{a_0 b_0}{d}$ .

b) Suppose  $k = 2t + 1$  ( $t = 1, 2, \dots$ ). Iterating the right-hand side of equations in

(9), we can write

$$\begin{aligned}
 x_{4tm+2m+2j+1} &= x_{2j-4t-1} \prod_{s=0}^m u_{2(2ts+s+j)+1} u_{2(2ts+s+j-t)} \\
 &+ \sum_{l=0}^{2m+1} \beta_{2(tl+j-t)+l} \prod_{s=l+1}^{2m+1} u_{2(ts+j-t)+s}
 \end{aligned} \tag{15}$$

$$\begin{aligned}
 x_{4tm+2m+2j+2} &= x_{2j-4t} \prod_{s=0}^m u_{2(2ts+s+j+1)} u_{2(2ts+s+j-t)+1} \\
 &+ \sum_{l=0}^{2m+1} \beta_{2(tl+j-t)+l+1} \prod_{s=l+1}^{2m+1} u_{2(ts+j-t)+s+1}
 \end{aligned} \tag{16}$$

$$\begin{aligned}
 y_{4tm+2m+2j+1} &= y_{2j-4t-1} \prod_{s=0}^m v_{2(2ts+s+j)+1} v_{2(2ts+s+j-t)} \\
 &+ \sum_{l=0}^{2m+1} \alpha_{2(tl+j-t)+l} \prod_{s=l+1}^{2m+1} v_{2(ts+j-t)+s}
 \end{aligned} \tag{17}$$

$$\begin{aligned}
 y_{4tm+2m+2j+2} &= y_{2j-4t} \prod_{s=0}^m v_{2(2ts+s+j+1)} v_{2(2ts+s+j-t)+1} \\
 &+ \sum_{l=0}^{2m+1} \alpha_{2(tl+j-t)+l+1} \prod_{s=l+1}^{2m+1} v_{2(ts+j-t)+s+1},
 \end{aligned} \tag{18}$$

where  $j \in \{t, t + 1, \dots, 3t\}$ . Using (8) in (15)-(18), we get

$$\begin{aligned}
 x_{4tm+2m+2j+1} &= x_{2j-4t-1} \prod_{s=0}^m d \left( \frac{a_0}{b_1} \right)^{(2t+1)s+j} \left( \frac{a_1}{b_0} \right)^{(2t+1)s+j-t} \\
 &+ \sum_{l=0}^m \beta_{2(2tl+j+l-t)} u_0^{m-l} u_1^{m-l+1} \\
 &\times \left( \frac{a_1}{b_0} \right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}} \left( \frac{a_0}{b_1} \right)^{j(m-l+1)+(2t+1)\frac{m(m+1)-l(l-1)}{2}} \\
 &+ \sum_{l=0}^m \beta_{2(2tl+j+l)+1} u_0^{m-l} u_1^{m-l} \left( \frac{a_1}{b_0} \right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l+1)}{2}} \\
 &\times \left( \frac{a_0}{b_1} \right)^{j(m-l)+(2t+1)\frac{m(m+1)-l(l+1)}{2}}
 \end{aligned}$$

$$x_{4tm+2m+2j+2} = x_{2j-4t} \prod_{s=0}^m d \left( \frac{a_0}{b_1} \right)^{(2t+1)s+j+1} \left( \frac{a_1}{b_0} \right)^{(2t+1)s+j-t}$$

$$\begin{aligned}
 & + \sum_{l=0}^m \beta_{2(2tl+j+l-t)+1} u_0^{m-l+1} u_1^{m-l} \\
 & \times \left(\frac{a_1}{b_0}\right)^{(j+1)(m+1-l)+(2t+1)\frac{m(m+1)-l(l-1)}{2}} \left(\frac{a_0}{b_1}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}} \\
 & + \sum_{l=0}^m \beta_{2(2tl+j+l+1)} u_0^{m-l} u_1^{m-l} \left(\frac{a_1}{b_0}\right)^{(j+1)(m-l)+(2t+1)\frac{m(m+1)-l(l+1)}{2}} \\
 & \times \left(\frac{a_0}{b_1}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}} \\
 \\
 y_{4tm+2m+2j+1} & = y_{2j-4t-1} \prod_{s=0}^m \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^{(2t+1)s+j} \left(\frac{b_1}{a_0}\right)^{(2t+1)s+j-t} \\
 & + \sum_{l=0}^m \alpha_{2(2tl+j+l-t)} v_0^{m-l} v_1^{m-l+1} \\
 & \times \left(\frac{b_1}{a_0}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}} \left(\frac{b_0}{a_1}\right)^{j(m-l+1)+(2t+1)\frac{m(m+1)-l(l-1)}{2}} \\
 & + \sum_{l=0}^m \alpha_{2(2tl+j+l+1)} v_0^{m-l} v_1^{m-l} \left(\frac{b_1}{a_0}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l+1)}{2}} \\
 & \times \left(\frac{b_0}{a_1}\right)^{j(m-l)+(2t+1)\frac{m(m+1)-l(l+1)}{2}} \\
 \\
 y_{4tm+2m+2j+2} & = y_{2j-4t} \prod_{s=0}^m \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^{(2t+1)s+j+1} \left(\frac{b_1}{a_0}\right)^{(2t+1)s+j-t} \\
 & + \sum_{l=0}^m \alpha_{2(2tl+j+l-t)+1} v_0^{m-l+1} v_1^{m-l} \left(\frac{b_1}{a_0}\right)^{(j+1)(m-l+1)+(2t+1)\frac{m(m+1)-l(l-1)}{2}} \\
 & \times \left(\frac{b_0}{a_1}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}} + \sum_{l=0}^m \alpha_{2(2tl+j+l+1)} v_0^{m-l} v_1^{m-l} \\
 & \times \left(\frac{b_1}{a_0}\right)^{(j+1)(m-l)+(2t+1)\frac{m(m+1)-l(l+1)}{2}} \left(\frac{b_0}{a_1}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}}
 \end{aligned}$$

c) Suppose  $k = (2t)$  ( $t = 1, 2, \dots$ ). Similarly, iterating the right-hand side of equations in (9), we have

$$x_{4tm+2j} = x_{2j-4t} \prod_{s=0}^m u_{2(2ts+j)} u_{2(2ts+j-t)}$$

$$+ \sum_{l=0}^{2m+1} \beta_{2(tl+j-t)} \prod_{s=l+1}^{2m+1} u_{2(ts+j-t)}, \tag{19}$$

$$\begin{aligned} x_{4tm+2j+1} &= x_{2j-4t+1} \prod_{s=0}^m u_{2(2ts+j)+1} u_{2(2ts+j-t)+1} \\ &+ \sum_{l=0}^{2m+1} \beta_{2(tl+j-t)+1} \prod_{s=l+1}^{2m+1} u_{2(ts+j-t)+1}, \end{aligned} \tag{20}$$

$$\begin{aligned} y_{4tm+2j} &= y_{2j-4t} \prod_{s=0}^m v_{2(2ts+j)} v_{2(2ts+j-t)} \\ &+ \sum_{l=0}^{2m+1} \alpha_{2(tl+j-t)} \prod_{s=l+1}^{2m+1} v_{2(ts+j-t)}, \end{aligned} \tag{21}$$

$$\begin{aligned} y_{4tm+2j+1} &= y_{2j-4t+1} \prod_{s=0}^m v_{2(2ts+j)+1} v_{2(2ts+j-t)+1} \\ &+ \sum_{l=0}^{2m+1} \alpha_{2(tl+j-t)+1} \prod_{s=l+1}^{2m+1} v_{2(ts+j-t)+1}, \end{aligned} \tag{22}$$

where  $j \in \{t, t + 1, \dots, 3t - 1\}$ . Using (8) in (19)-(22), we get

$$\begin{aligned} x_{4tm+2j} &= x_{2j-4t} \prod_{s=0}^m u_0^2 \left(\frac{a_1}{b_0}\right)^{4ts+2j-t} + \sum_{l=0}^{2m+1} \beta_{2(tl+j-t)} u_0^{2m-l+1} \\ &\times \left(\frac{a_1}{b_0}\right)^{j(2m-l+1)+t \frac{2m(2m+1)-l(l-1)}{2}} \\ x_{4tm+2j+1} &= x_{2j-4t+1} \prod_{s=0}^m u_1^2 \left(\frac{a_0}{b_1}\right)^{4ts+2j-t} + \sum_{l=0}^{2m+1} \beta_{2(tl+j-t)+1} u_1^{2m-l+1} \\ &\times \left(\frac{a_0}{b_1}\right)^{j(2m-l+1)+t \frac{2m(2m+1)-l(l-1)}{2}} \\ y_{4tm+2j} &= y_{2j-4t} \prod_{s=0}^m v_0^2 \left(\frac{b_1}{a_0}\right)^{4ts+2j-t} + \sum_{l=0}^{2m+1} \alpha_{2(tl+j-t)} v_0^{2m-l+1} \\ &\times \left(\frac{b_1}{a_0}\right)^{j(2m-l+1)+t \frac{2m(2m+1)-l(l-1)}{2}} \\ y_{4tm+2j+1} &= y_{2j-4t+1} \prod_{s=0}^m v_1^2 \left(\frac{b_0}{a_1}\right)^{4ts+2j-t} + \sum_{l=0}^{2m+1} \alpha_{2(tl+j-t)+1} v_1^{2m-l+1} \end{aligned}$$

$$\times \left(\frac{b_0}{a_1}\right)^{j(2m-l+1)+t\frac{2m(2m+1)-l(l-1)}{2}},$$

where  $u_0 = \frac{x_0-\beta_0}{x-k}$ ,  $u_1 = \frac{x_1-\beta_1}{x_1-k}$ ,  $v_0 = \frac{y_0-\alpha_0}{y-k}$ ,  $v_1 = \frac{y_1-\alpha_1}{y_1-k}$ ,  $d = u_0u_1$  and  $v_0v_1 = \frac{a_0b_0}{d}$ .

The previous computations prove the next theorem.

**Theorem 3.** *Let  $\{(x_n, y_n)\}_{n \geq -k}$  be a well-defined solution of system (3). Then, we get the next formulas:*

(a) *If  $k = 1$ , for all  $n \in N_0$ , then we get*

$$\begin{aligned} x_{2n-1} &= x_{-1} \prod_{t=0}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^t + \sum_{r=0}^{n-1} \beta_0 u_1 \left(\frac{a_0}{b_1}\right)^r \prod_{t=r+1}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^t \\ &+ \sum_{r=0}^{n-1} \beta_1 \prod_{t=r+1}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^t, \\ x_{2n} &= x_0 \prod_{t=0}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^{t+1} + \sum_{r=0}^{n-1} \beta_0 \prod_{t=r+1}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^{t+1} \\ &+ \sum_{r=0}^{n-1} \beta_1 u_0 \left(\frac{a_1}{b_0}\right)^{r+1} \prod_{t=r+1}^{n-1} d \left(\frac{a_0}{b_1}\right)^t \left(\frac{a_1}{b_0}\right)^{t+1}, \\ y_{2n-1} &= y_{-1} \prod_{t=0}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^t + \sum_{r=0}^{n-1} \alpha_0 v_1 \left(\frac{b_0}{a_1}\right)^r \prod_{t=r+1}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^t \\ &+ \sum_{r=0}^{n-1} \alpha_1 \prod_{t=r+1}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^t, \\ y_{2n} &= y_0 \prod_{t=0}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^{t+1} + \sum_{r=0}^{n-1} \alpha_0 \prod_{t=r+1}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^{t+1} \\ &+ \sum_{r=0}^{n-1} \alpha_1 v_0 \left(\frac{b_1}{a_0}\right)^{r+1} \prod_{t=r+1}^{n-1} \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^t \left(\frac{b_1}{a_0}\right)^{t+1}. \end{aligned}$$

b) *If  $k = 2t + 1$  ( $t = 1, 2, \dots$ ), for all  $m \in N_0$ , we have*

$$\begin{aligned} x_{4tm+2m+2j+1} &= x_{2j-4t-1} \prod_{s=0}^m d \left(\frac{a_0}{b_1}\right)^{(2t+1)s+j} \left(\frac{a_1}{b_0}\right)^{(2t+1)s+j-t} \\ &+ \sum_{l=0}^m \beta_{2(2tl+j+l-t)} u_0^{m-l} u_1^{m-l+1} \left(\frac{a_1}{b_0}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}} \end{aligned}$$



$$\begin{aligned}
 & \times \left(\frac{a_0}{b_1}\right)^{j(m-l+1)+(2t+1)\frac{m(m+1)-l(l-1)}{2}} \\
 & + \sum_{l=0}^m \beta_{2(2tl+j+l)+1} u_0^{m-l} u_1^{m-l} \left(\frac{a_1}{b_0}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l+1)}{2}} \\
 & \times \left(\frac{a_0}{b_1}\right)^{j(m-l)+(2t+1)\frac{m(m+1)-l(l+1)}{2}}
 \end{aligned}$$

$$\begin{aligned}
 x_{4tm+2m+2j+2} &= x_{2j-4t} \prod_{s=0}^m d \left(\frac{a_0}{b_1}\right)^{(2t+1)s+j+1} \left(\frac{a_1}{b_0}\right)^{(2t+1)s+j-t} \\
 & + \sum_{l=0}^m \beta_{2(2tl+j+l-t)+1} u_0^{m-l+1} u_1^{m-l} \\
 & \times \left(\frac{a_1}{b_0}\right)^{(j+1)(m-l+1)+(2t+1)\frac{m(m+1)-l(l-1)}{2}} \left(\frac{a_0}{b_1}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}} \\
 & + \sum_{l=0}^m \beta_{2(2tl+j+l+1)} u_0^{m-l} u_1^{m-l} \left(\frac{a_1}{b_0}\right)^{(j+1)(m-l)+(2t+1)\frac{m(m+1)-l(l+1)}{2}} \\
 & \times \left(\frac{a_0}{b_1}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}}
 \end{aligned}$$

$$\begin{aligned}
 y_{4tm+2m+2j+1} &= y_{2j-4t-1} \prod_{s=0}^m \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^{(2t+1)s+j} \left(\frac{b_1}{a_0}\right)^{(2t+1)s+j-t} \\
 & + \sum_{l=0}^m \alpha_{2(2tl+j+l-t)} v_0^{m-l} v_1^{m-l+1} \\
 & \times \left(\frac{b_1}{a_0}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}} \left(\frac{b_0}{a_1}\right)^{j(m-l+1)+(2t+1)\frac{m(m+1)-l(l-1)}{2}} \\
 & + \sum_{l=0}^m \alpha_{2(2tl+j+l)+1} v_0^{m-l} v_1^{m-l} \left(\frac{b_1}{a_0}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l+1)}{2}} \\
 & \times \left(\frac{b_0}{a_1}\right)^{j(m-l)+(2t+1)\frac{m(m+1)-l(l+1)}{2}}
 \end{aligned}$$

$$y_{4tm+2m+2j+2} = y_{2j-4t} \prod_{s=0}^m \frac{a_0 b_0}{d} \left(\frac{b_0}{a_1}\right)^{(2t+1)s+j+1} \left(\frac{b_1}{a_0}\right)^{(2t+1)s+j-t}$$

$$\begin{aligned}
 & + \sum_{l=0}^m \alpha_{2(2tl+j+l-t)+1} v_0^{m-l+1} v_1^{m-l} \left(\frac{b_1}{a_0}\right)^{(j+1)(m-l+1)+(2t+1)\frac{m(m+1)-l(l-1)}{2}} \\
 & \times \left(\frac{b_0}{a_1}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}} + \sum_{l=0}^m \alpha_{2(2tl+j+l+1)} v_0^{m-l} v_1^{m-l} \\
 & \times \left(\frac{b_1}{a_0}\right)^{(j+1)(m-l)+(2t+1)\frac{m(m+1)-l(l+1)}{2}} \left(\frac{b_0}{a_1}\right)^{(t+j+1)(m-l)+(2t+1)\frac{m(m-1)-l(l-1)}{2}}
 \end{aligned}$$

c) If  $k = 2t$  ( $t = 1, 2, \dots$ ), for all  $m \in N_0$ , we have

$$\begin{aligned}
 x_{4tm+2j} &= x_{2j-4t} \prod_{s=0}^m u_0^2 \left(\frac{a_1}{b_0}\right)^{4ts+2j-t} + \sum_{l=0}^{2m+1} \beta_{2(tl+j-t)} u_0^{2m-l+1} \\
 & \times \left(\frac{a_1}{b_0}\right)^{j(2m-l+1)+t\frac{2m(2m+1)-l(l-1)}{2}} \\
 x_{4tm+2j+1} &= x_{2j-4t+1} \prod_{s=0}^m u_1^2 \left(\frac{a_0}{b_1}\right)^{4ts+2j-t} + \sum_{l=0}^{2m+1} \beta_{2(tl+j-t)+1} u_1^{2m-l+1} \\
 & \times \left(\frac{a_0}{b_1}\right)^{j(2m-l+1)+t\frac{2m(2m+1)-l(l-1)}{2}} \\
 y_{4tm+2j} &= y_{2j-4t} \prod_{s=0}^m v_0^2 \left(\frac{b_1}{a_0}\right)^{4ts+2j-t} + \sum_{l=0}^{2m+1} \alpha_{2(tl+j-t)} v_0^{2m-l+1} \\
 & \times \left(\frac{b_1}{a_0}\right)^{j(2m-l+1)+t\frac{2m(2m+1)-l(l-1)}{2}} \\
 y_{4tm+2j+1} &= y_{2j-4t+1} \prod_{s=0}^m v_1^2 \left(\frac{b_0}{a_1}\right)^{4ts+2j-t} + \sum_{l=0}^{2m+1} \alpha_{2(tl+j-t)+1} v_1^{2m-l+1} \\
 & \times \left(\frac{b_0}{a_1}\right)^{j(2m-l+1)+t\frac{2m(2m+1)-l(l-1)}{2}},
 \end{aligned}$$

where  $u_0 = \frac{x_0-\beta_0}{x_{-k}}$ ,  $u_1 = \frac{x_1-\beta_1}{x_{1-k}}$ ,  $v_0 = \frac{y_0-\alpha_0}{y_{-k}}$ ,  $v_1 = \frac{y_1-\alpha_1}{y_{1-k}}$ ,  $d = u_0 u_1$  and  $v_0 v_1 = \frac{a_0 b_0}{d}$ .

### 3. THE CASE $a_0 = b_1$ AND $a_1 = b_0$

The aim of in this section is to investigate the asymptotic behavior and periodicity of well-defined solutions of system (3) in the case  $a_0 = b_1$  and  $a_1 = b_0$ . The following corollary gives us the solutions of system (3) when  $a_0 = b_1$  and  $a_1 = b_0$ .

**Corollary 4.** *Suppose that  $\{(x_n, y_n)\}_{n \geq -k}$  is a well-defined solution of system (3) when  $a_0 = b_1$  and  $a_1 = b_0$ . Then, the solutions of system (3) are given by:*

a) If  $k = 1$ , for all  $m \in N_0$ , then we get

$$x_{2m-1} = \begin{cases} x_{-1} + (\beta_0 u_1 + \beta_1) m & \text{if } d = 1, \\ x_{-1} d^m + (\beta_0 u_1 + \beta_1) \frac{d^m - 1}{d - 1} & \text{if } d \neq 1, \end{cases} \quad (23)$$

$$x_{2m} = \begin{cases} x_0 + (\beta_1 u_0 + \beta_0) m & \text{if } d = 1, \\ x_0 d^m + (\beta_1 u_0 + \beta_0) \frac{d^m - 1}{d - 1} & \text{if } d \neq 1, \end{cases} \quad (24)$$

$$y_{2m-1} = \begin{cases} y_{-1} + (\alpha_0 v_1 + \alpha_1) m & \text{if } a_0 b_0 = d, \\ y_{-1} \left(\frac{a_0 b_0}{d}\right)^m + (\alpha_0 v_1 + \alpha_1) \frac{\left(\frac{a_0 b_0}{d}\right)^m - 1}{\frac{a_0 b_0}{d} - 1} & \text{if } a_0 b_0 \neq d, \end{cases} \quad (25)$$

$$y_{2m} = \begin{cases} y_0 + (\alpha_1 v_0 + \alpha_0) m & \text{if } a_0 b_0 = d, \\ y_0 \left(\frac{a_0 b_0}{d}\right)^m + (\alpha_1 v_0 + \alpha_0) \frac{\left(\frac{a_0 b_0}{d}\right)^m - 1}{\frac{a_0 b_0}{d} - 1} & \text{if } a_0 b_0 \neq d, \end{cases} \quad (26)$$

b) If  $k = 2t + 1$  ( $t = 1, 2, \dots$ ), for all  $m \in N_0$  and  $j \in \{t, t + 1, \dots, 3t\}$ , we have

$$x_{4tm+2m+2j+1} = \begin{cases} x_{2j-4t-1} + (\beta_0 u_1 + \beta_1) (m + 1) & \text{if } d = 1, \\ x_{2j-4t-1} d^{m+1} + (\beta_0 u_1 + \beta_1) \left(\frac{d^{m+1} - 1}{d - 1}\right) & \text{if } d \neq 1, \end{cases} \quad (27)$$

$$x_{4tm+2m+2j+2} = \begin{cases} x_{2j-4t} + (\beta_1 u_0 + \beta_0) (m + 1) & \text{if } d = 1, \\ x_{2j-4t} d^{m+1} + (\beta_1 u_0 + \beta_0) \left(\frac{d^{m+1} - 1}{d - 1}\right) & \text{if } d \neq 1, \end{cases} \quad (28)$$

$$y_{4tm+2m+2j+1} = \begin{cases} y_{2j-4t-1} + (\alpha_0 v_1 + \alpha_1) (m + 1) & \text{if } d = a_0 b_0, \\ y_{2j-4t-1} \left(\frac{a_0 b_0}{d}\right)^{m+1} + (\alpha_0 v_1 + \alpha_1) \left(\frac{\left(\frac{a_0 b_0}{d}\right)^{m+1} - 1}{\frac{a_0 b_0}{d} - 1}\right) & \text{if } d \neq a_0 b_0, \end{cases} \quad (29)$$

$$y_{4tm+2m+2j+2} = \begin{cases} y_{2j-4t} + (\alpha_1 v_0 + \alpha_0) (m + 1) & \text{if } d = a_0 b_0 \\ y_{2j-4t} \left(\frac{a_0 b_0}{d}\right)^{m+1} + (\alpha_1 v_0 + \alpha_0) \left(\frac{\left(\frac{a_0 b_0}{d}\right)^{m+1} - 1}{\frac{a_0 b_0}{d} - 1}\right) & \text{if } d \neq a_0 b_0, \end{cases} \quad (30)$$

c) If  $k = 2t$  ( $t = 1, 2, \dots$ ), for all  $m \in N_0$  and  $j \in \{t, t + 1, \dots, 3t - 1\}$ , we have

$$x_{4tm+2j+1} = \begin{cases} x_{2j-4t+1} + \beta_1 (2m + 2) & \text{if } u_1 = 1, \\ x_{2j-4t+1} u_1^{2m+2} + \beta_1 \left(\frac{u_1^{2m+2} - 1}{u_1 - 1}\right) & \text{if } u_1 \neq 1, \end{cases} \quad (31)$$

$$x_{4tm+2j} = \begin{cases} x_{2j-4t} + \beta_0 (2m + 2) & \text{if } u_0 = 1, \\ x_{2j-4t} (u_0)^{2m+2} + \beta_0 \left(\frac{(u_0)^{2m+2} - 1}{u_0 - 1}\right) & \text{if } u_0 \neq 1, \end{cases} \quad (32)$$

$$y_{4tm+2j+1} = \begin{cases} y_{2j-4t+1} + \alpha_1 (2m + 2) & \text{if } v_1 = 1, \\ y_{2j-4t+1} (v_1)^{2m+2} + \alpha_1 \left(\frac{(v_1)^{2m+2} - 1}{v_1 - 1}\right) & \text{if } v_1 \neq 1, \end{cases} \quad (33)$$

$$y_{4tm+2j} = \begin{cases} y_{2j-4t} + \alpha_0 (2m + 2) & \text{if } v_0 = 1, \\ y_{2j-4t} (v_0)^{2m+2} + \alpha_0 \left( \frac{(v_0)^{2m+2}-1}{v_0-1} \right) & \text{if } v_0 \neq 1. \end{cases} \tag{34}$$

where  $u_0 = \frac{x_0-\beta_0}{x_{-k}}$ ,  $u_1 = \frac{x_1-\beta_1}{x_{1-k}}$ ,  $v_0 = \frac{y_0-\alpha_0}{y_{-k}}$ ,  $v_1 = \frac{y_1-\alpha_1}{y_{1-k}}$ ,  $d = u_0u_1$  and  $v_0v_1 = \frac{a_0b_0}{d}$ .

The next theorem gives the limiting properties of solutions of system (3) in the case  $a_0 = b_1$  and  $a_1 = b_0$ .

**Theorem 5.** *Suppose that  $\{(x_n, y_n)\}_{n \geq -k}$  is a well-defined solution of system (3) with  $a_0 = b_1$ ,  $a_1 = b_0$ . Then, the next statements are true.*

- (1) Let  $k = 1$ .
  - (a) If  $(d - 1)x_{-1} + (\beta_0u_1 + \beta_1) \neq 0$ , then we get

$$\lim_{n \rightarrow \infty} |x_{2n-1}| = \begin{cases} \left| \frac{\beta_0u_1 + \beta_1}{d-1} \right| & \text{if } |d| < 1, \\ \infty & \text{if } |d| > 1. \end{cases}$$

Otherwise, if  $(d - 1)x_{-1} + (\beta_0u_1 + \beta_1) = 0$  and  $d \neq 1$ , then  $x_{2n-1} = x_{-1}$  for all  $n \in N_0$ .

- (b) If  $\beta_0u_1 + \beta_1 \neq 0$  and  $d = 1$ , then  $|x_{2n-1}| \rightarrow \infty$ , as  $n \rightarrow \infty$ . Otherwise, if  $\beta_0u_1 + \beta_1 = 0$  and  $d = 1$ , then  $x_{2n-1} = x_{-1}$ , for all  $n \in N_0$ .
- (c) If  $(d - 1)x_0 + (\beta_1u_0 + \beta_0) \neq 0$ , then we have

$$\lim_{n \rightarrow \infty} |x_{2n}| = \begin{cases} \left| \frac{\beta_1u_0 + \beta_0}{d-1} \right| & \text{if } |d| < 1, \\ \infty & \text{if } |d| > 1. \end{cases}$$

Otherwise, if  $(d - 1)x_0 + (\beta_1u_0 + \beta_0) = 0$  and  $d \neq 1$ , then  $x_{2n} = x_0$  for all  $n \in N_0$ .

- (d) If  $\beta_1u_0 + \beta_0 \neq 0$  and  $d = 1$ , then  $|x_{2n}| \rightarrow \infty$ , as  $n \rightarrow \infty$ . Otherwise, if  $\beta_1u_0 + \beta_0 = 0$  and  $d = 1$ , then  $x_{2n} = x_0$ , for all  $n \in N_0$ .
- (e) If  $\left(\frac{a_0b_0}{d} - 1\right)y_{-1} + (\alpha_0v_1 + \alpha_1) \neq 0$ , then we get

$$\lim_{n \rightarrow \infty} |y_{2n-1}| = \begin{cases} \left| \frac{(\alpha_0v_1 + \alpha_1)d}{d - a_0b_0} \right| & \text{if } |d| > a_0b_0, \\ \infty & \text{if } |d| < a_0b_0. \end{cases}$$

Otherwise, if  $\left(\frac{a_0b_0}{d} - 1\right)y_{-1} + (\alpha_0v_1 + \alpha_1) = 0$  and  $d \neq a_0b_0$ , then  $y_{2n-1} = y_{-1}$  for all  $n \in N_0$ .

- (f) If  $\alpha_0v_1 + \alpha_1 \neq 0$  and  $d = a_0b_0$ , then  $|y_{2n-1}| \rightarrow \infty$ , as  $n \rightarrow \infty$ . Otherwise, if  $\alpha_0v_1 + \alpha_1 = 0$  and  $d = a_0b_0$ , then  $y_{2n-1} = y_{-1}$ , for all  $n \in N_0$ .
- (g) If  $\left(\frac{a_0b_0}{d} - 1\right)y_0 + (\alpha_1v_0 + \alpha_0) \neq 0$ , then we get

$$\lim_{n \rightarrow \infty} |y_{2n}| = \begin{cases} \left| \frac{(\alpha_1v_0 + \alpha_0)d}{d - a_0b_0} \right| & \text{if } |d| > a_0b_0 \\ \infty & \text{if } |d| < a_0b_0 \end{cases}$$

Otherwise, if  $(\frac{a_0 b_0}{d} - 1) y_0 + (\alpha_1 v_0 + \alpha_0) = 0$  and  $d \neq a_0 b_0$ , then  $y_{2n} = y_0$  for all  $n \in N_0$ .

(h) If  $\alpha_1 v_0 + \alpha_0 \neq 0$  and  $d = a_0 b_0$ , then  $|y_{2n}| \rightarrow \infty$ , as  $n \rightarrow \infty$ . Otherwise, if  $\alpha_1 v_0 + \alpha_0 = 0$  and  $d = a_0 b_0$ , then  $y_{2n} = y_0$ , for all  $n \in N_0$ .

(2) Let  $k = 2t + 1$  ( $t = 1, 2, \dots$ ) and  $j \in \{t, t + 1, \dots, 3t\}$ .

(a) If  $(d - 1) x_{2j-4t-1} + (\beta_0 u_1 + \beta_1) \neq 0$ , then we have

$$\lim_{m \rightarrow \infty} |x_{4tm+2m+2j+1}| = \begin{cases} \left| \frac{\beta_0 u_1 + \beta_1}{d-1} \right| & \text{if } |d| < 1, \\ \infty & \text{if } |d| > 1. \end{cases}$$

Otherwise, if  $(d - 1) x_{2j-4t-1} + (\beta_0 u_1 + \beta_1) = 0$  and  $d \neq 1$ , then  $x_{4tm+2m+2j+1} = x_{2j-4t-1}$  for all  $m \in N_0$ .

(b) If  $\beta_0 u_1 + \beta_1 \neq 0$  and  $d = 1$ , then  $|x_{4tm+2m+2j+1}| \rightarrow \infty$ , as  $m \rightarrow \infty$ . Otherwise, if  $\beta_0 u_1 + \beta_1 = 0$  and  $d = 1$ , then  $x_{4tm+2m+2j+1} = x_{2j-4t-1}$ , for all  $m \in N_0$ .

(c) If  $(d - 1) x_{2j-4t} + (\beta_1 u_0 + \beta_0) \neq 0$ , then we get

$$\lim_{m \rightarrow \infty} |x_{4tm+2m+2j+2}| = \begin{cases} \left| \frac{\beta_1 u_0 + \beta_0}{d-1} \right| & \text{if } |d| < 1, \\ \infty & \text{if } |d| > 1. \end{cases}$$

Otherwise, if  $(d - 1) x_{2j-4t} + (\beta_1 u_0 + \beta_0) = 0$  and  $d \neq 1$ , then  $x_{4tm+2m+2j+2} = x_{2j-4t}$  for all  $m \in N_0$ .

(d) If  $\beta_1 u_0 + \beta_0 \neq 0$  and  $d = 1$ , then  $|x_{4tm+2m+2j+2}| \rightarrow \infty$ , as  $m \rightarrow \infty$ . Otherwise, if  $\beta_1 u_0 + \beta_0 = 0$  and  $d = 1$ , then  $x_{4tm+2m+2j+2} = x_{2j-4t}$ , for all  $m \in N_0$ .

(e) If  $(\frac{a_0 b_0}{d} - 1) y_{2j-4t-1} + (\alpha_0 v_1 + \alpha_1) \neq 0$ , then we have

$$\lim_{m \rightarrow \infty} |y_{4tm+2m+2j+1}| = \begin{cases} \left| \frac{(\alpha_0 v_1 + \alpha_1)d}{d - a_0 b_0} \right| & \text{if } |d| > a_0 b_0, \\ \infty & \text{if } |d| < a_0 b_0. \end{cases}$$

Otherwise, if  $(\frac{a_0 b_0}{d} - 1) y_{2j-4t-1} + (\alpha_0 v_1 + \alpha_1) = 0$  and  $d \neq a_0 b_0$ , then  $y_{4tm+2m+2j+1} = y_{2j-4t-1}$  for all  $m \in N_0$ .

(f) If  $\alpha_0 v_1 + \alpha_1 \neq 0$  and  $d = a_0 b_0$ , then  $|y_{4tm+2m+2j+1}| \rightarrow \infty$ , as  $m \rightarrow \infty$ . Otherwise, if  $\alpha_0 v_1 + \alpha_1 = 0$  and  $d = a_0 b_0$ , then  $y_{4tm+2m+2j+1} = y_{2j-4t-1}$ , for all  $m \in N_0$ .

(g) If  $(\frac{a_0 b_0}{d} - 1) y_{2j-4t} + (\alpha_1 v_0 + \alpha_0) \neq 0$ , then we get

$$\lim_{m \rightarrow \infty} |y_{4tm+2m+2j+2}| = \begin{cases} \left| \frac{(\alpha_1 v_0 + \alpha_0)d}{d - a_0 b_0} \right| & \text{if } |d| > a_0 b_0, \\ \infty & \text{if } |d| < a_0 b_0. \end{cases}$$

Otherwise, if  $(\frac{a_0 b_0}{d} - 1) y_{2j-4t} + (\alpha_1 v_0 + \alpha_0) = 0$  and  $d \neq a_0 b_0$ , then  $y_{4tm+2m+2j+2} = y_{2j-4t}$  for all  $m \in N_0$ .

- (h) If  $\alpha_1 v_0 + \alpha_0 \neq 0$  and  $d = a_0 b_0$ , then  $|y_{4tm+2m+2j+2}| \rightarrow \infty$ , as  $m \rightarrow \infty$ .  
 Otherwise, if  $\alpha_1 v_0 + \alpha_0 = 0$  and  $d = a_0 b_0$ , then  $y_{4tm+2m+2j+2} = y_{2j-4t}$ ,  
 for all  $m \in N_0$ .

- (3) Let  $k = 2t$  ( $t = 1, 2, \dots$ ) and  $j \in \{t, t+1, \dots, 3t-1\}$ .

- (a) If  $(u_1 - 1)x_{2j-4t+1} + \beta_1 \neq 0$ , then we have

$$\lim_{m \rightarrow \infty} |x_{4tm+2j+1}| = \begin{cases} \left| \frac{\beta_1}{u_1-1} \right| & \text{if } |u_1| < 1, \\ \infty & \text{if } |u_1| > 1. \end{cases}$$

Otherwise, if  $(u_1 - 1)x_{2j-4t+1} + \beta_1 = 0$  and  $u_1 \neq 1$ , then  $x_{4tm+2j+1} = x_{2j-4t+1}$  for all  $m \in N_0$ .

- (b) If  $\beta_1 \neq 0$  and  $u_1 = 1$ , then  $|x_{4tm+2j+1}| \rightarrow \infty$ , as  $m \rightarrow \infty$ . Otherwise,  
 if  $\beta_1 = 0$  and  $u_1 = 1$ , then  $x_{4tm+2j+1} = x_{2j-4t+1}$ , for all  $m \in N_0$ .  
 (c) If  $(u_0 - 1)x_{2j-4t} + \beta_0 \neq 0$ , then we get

$$\lim_{m \rightarrow \infty} |x_{4tm+2j}| = \begin{cases} \left| \frac{\beta_0}{u_0-1} \right| & \text{if } |u_0| < 1, \\ \infty & \text{if } |u_0| > 1. \end{cases}$$

Otherwise, if  $(u_0 - 1)x_{2j-4t} + \beta_0 = 0$  and  $u_0 \neq 1$ , then  $x_{4tm+2j} = x_{2j-4t}$  for all  $m \in N_0$ .

- (d) If  $\beta_0 \neq 0$  and  $u_0 = 1$ , then  $|x_{4tm+2j}| \rightarrow \infty$ , as  $m \rightarrow \infty$ . Otherwise,  
 if  $\beta_0 = 0$  and  $u_0 = 1$ , then  $x_{4tm+2j} = x_{2j-4t}$ , for all  $m \in N_0$ .  
 (e) If  $(v_1 - 1)y_{2j-4t+1} + \alpha_1 \neq 0$ , then we have

$$\lim_{m \rightarrow \infty} |y_{4tm+2j+1}| = \begin{cases} \left| \frac{\alpha_1}{v_1-1} \right| & \text{if } |v_1| < 1, \\ \infty & \text{if } |v_1| > 1. \end{cases}$$

Otherwise, if  $(v_1 - 1)y_{2j-4t+1} + \alpha_1 = 0$  and  $v_1 \neq 1$ , then  $y_{4tm+2j+1} = y_{2j-4t+1}$  for all  $m \in N_0$ .

- (f) If  $\alpha_1 \neq 0$  and  $v_1 = 1$ , then  $|y_{4tm+2j+1}| \rightarrow \infty$ , as  $m \rightarrow \infty$ . Otherwise,  
 if  $\alpha_1 = 0$  and  $v_1 = 1$ , then  $y_{4tm+2j+1} = y_{2j-4t+1}$ , for all  $m \in N_0$ .  
 (g) If  $(v_0 - 1)y_{2j-4t} + \alpha_0 \neq 0$ , then we get

$$\lim_{m \rightarrow \infty} |y_{4tm+2j}| = \begin{cases} \left| \frac{\alpha_0}{v_0-1} \right| & \text{if } |v_0| < 1, \\ \infty & \text{if } |v_0| > 1. \end{cases}$$

Otherwise, if  $(v_0 - 1)y_{2j-4t} + \alpha_0 = 0$  and  $v_0 \neq 1$ , then  $y_{4tm+2j} = y_{2j-4t}$  for all  $m \in N_0$ .

- (h) If  $\alpha_0 \neq 0$  and  $v_0 = 1$ , then  $|y_{4tm+2j}| \rightarrow \infty$ , as  $m \rightarrow \infty$ . Otherwise,  
 if  $\alpha_0 = 0$  and  $v_0 = 1$ , then  $y_{4tm+2j} = y_{2j-4t}$ , for all  $m \in N_0$ .

*Proof.* We will only prove the items (a) and (b) for  $k = 2t + 1$  ( $t = 1, 2, \dots$ ) and  $j \in \{t, t+1, \dots, 3t\}$  since the other cases can be proved similarly.

- (a) Assume that  $(d - 1)x_{2j-4t-1} + (\beta_0 u_1 + \beta_1) \neq 0$ . It is easy to see from Corollary 4 that  $x_{4tm+2m+2j+1} \neq 0$ . Clearly, if  $|d| < 1$ , then  $|d|^{m+1} \rightarrow 0$

as  $m \rightarrow \infty$ . On the other hand, if  $|d| > 1$ , then  $|d|^{m+1} \rightarrow \infty$  as  $m \rightarrow \infty$ . From Eq. (27), we get

$$\begin{aligned} \lim_{m \rightarrow \infty} |x_{4tm+2m+2j+1}| &= \lim_{m \rightarrow \infty} \left| \frac{(d-1)x_{2j-4t-1} + (\beta_0 u_1 + \beta_1)}{d-1} d^{m+1} \right. \\ &\quad \left. + \left( \frac{(\beta_0 u_1 + \beta_1)}{1-d} \right) \right| \\ &= \left| \frac{(d-1)x_{2j-4t-1} + (\beta_0 u_1 + \beta_1)}{d-1} \right| \lim_{m \rightarrow \infty} d^{m+1} \\ &\quad + \left| \left( \frac{(\beta_0 u_1 + \beta_1)}{1-d} \right) \right| \\ &= \begin{cases} \left| \frac{(\beta_0 u_1 + \beta_1)}{d-1} \right| & \text{if } |d| < 1 \\ \infty & \text{if } |d| > 1 \end{cases}. \end{aligned}$$

Now on the other hand  $(d-1)x_{2j-4t-1} + (\beta_0 u_1 + \beta_1) = 0$  and  $d \neq 1$ . Then we have

$$\begin{aligned} x_{4tm+2m+2j+1} &= x_{2j-4t-1} d^{m+1} + (\beta_0 u_1 + \beta_1) \left( \frac{d^{m+1} - 1}{d-1} \right) \\ &= x_{2j-4t-1} d^{m+1} + \left( \frac{d^{m+1} - 1}{d-1} \right) (-(d-1)x_{2j-4t-1}) \\ &= x_{2j-4t-1} d^{m+1} - (d^{m+1} - 1)x_{2j-4t-1} \\ &= x_{2j-4t-1}, \end{aligned}$$

which completes the proof of (a).

(b) Let  $d = 1$ . If  $\beta_0 u_1 + \beta_1 \neq 0$ , then from Eq. (27) we get

$$x_{4tm+2m+2j+1} = x_{2j-4t-1} + (\beta_0 u_1 + \beta_1)(m+1) \neq 0.$$

Letting  $m \rightarrow \infty$  in above equations implies that  $|x_{4tm+2m+2j+1}| \rightarrow \infty$ . On the other hand, If  $\beta_0 u_1 + \beta_1 = 0$ , then obviously, for all  $m \in N_0$ ,

$$x_{4tm+2m+2j+1} = x_{2j-4t-1} + (\beta_0 u_1 + \beta_1)(m+1) = x_{2j-4t-1} + 0 = x_{2j-4t-1},$$

which completes the proof of (b). □

**Corollary 6.** Suppose that  $\{(x_n, y_n)\}_{n \geq -k}$  is a well-defined solution of system (3) with  $a_0 = b_1, a_1 = b_0, u_0 u_1 = d, v_0 v_1 = \frac{a_0 b_0}{d}$ . Then, the next statements are true.

- (1) Let  $k = 1$ .

(a) If  $d = -1$ , then for all  $m \in N_0$ , we get

$$\begin{cases} x_{4m-1} &= x_{-1}, \\ x_{4m} &= x_0, \\ x_{4m+1} &= -x_{-1} + (\beta_0 u_1 + \beta_1), \\ x_{4m+2} &= -x_0 + (\beta_1 u_0 + \beta_0). \end{cases}$$

(b) If  $d = -a_0 b_0$ , then for all  $m \in N_0$ , we have

$$\begin{cases} y_{4m-1} &= y_{-1}, \\ y_{4m} &= y_0, \\ y_{4m+1} &= -y_{-1} + (\alpha_0 v_1 + \alpha_1), \\ y_{4m+2} &= -y_0 + (\alpha_1 v_0 + \alpha_0). \end{cases}$$

(c) If  $d = 1, u_1 = u_0 = 1, \beta_0 + \beta_1 = 0$ , then for all  $m \in N_0$ , we get

$$\begin{cases} x_{2m-1} &= x_{-1}, \\ x_{2m} &= x_0. \end{cases}$$

(d) If  $d = a_0 b_0, v_1 = v_0 = 1, \alpha_0 + \alpha_1 = 0$ , then for all  $m \in N_0$ , we have

$$\begin{cases} y_{2m-1} &= y_{-1}, \\ y_{2m} &= y_0. \end{cases}$$

(2) Let  $k = 2t + 1, (t = 1, 2, \dots)$  and  $j \in \{t, t + 1, \dots, 3t\}$ .

(a) If  $d = -1$ , then for all  $m \in N_0$ , we have

$$\begin{cases} x_{8tm+4m+2j+1} &= -x_{2j-4t-1} + (\beta_0 u_1 + \beta_1), \\ x_{8tm+4m+4t+2j+3} &= x_{2j-4t-1}, \\ x_{8tm+4m+2j+2} &= -x_{2j-4t} + (\beta_1 u_0 + \beta_0), \\ x_{8tm+4m+4t+2j+4} &= x_{2j-4t}. \end{cases}$$

(b) If  $d = -a_0 b_0$ , then for all  $m \in N_0$ , we get

$$\begin{cases} y_{8tm+4m+2j+1} &= -y_{2j-4t-1} + (\alpha_0 v_1 + \alpha_1), \\ y_{8tm+4m+4t+2j+3} &= y_{2j-4t-1}, \\ y_{8tm+4m+2j+2} &= -y_{2j-4t} + (\alpha_1 v_0 + \alpha_0), \\ y_{8tm+4m+4t+2j+4} &= y_{2j-4t}. \end{cases}$$

(c) If  $d = 1, u_1 = u_0 = 1, \beta_0 + \beta_1 = 0$  then for all  $m \in N_0$ , we have

$$\begin{cases} x_{4tm+2m+2j+1} &= x_{2j-4t-1}, \\ x_{4tm+2m+2j+2} &= x_{2j-4t}. \end{cases}$$



(d) If  $d = a_0 b_0$ ,  $v_1 = v_0 = 1$ ,  $\alpha_0 + \alpha_1 = 0$  then for all  $m \in N_0$ , we get

$$\begin{cases} y_{4tm+2m+2j+1} &= y_{2j-4t-1}, \\ y_{4tm+2m+2j+2} &= y_{2j-4t}. \end{cases}$$

(3) Let  $k = 2t$ , ( $t = 1, 2, \dots$ ) and  $j \in \{t, t+1, \dots, 3t-1\}$ . If  $u_1 = u_0 = v_1 = v_0 = -1$  or  $u_1 = u_0 = v_1 = v_0 = 1$  and  $\beta_1 = \beta_0 = \alpha_1 = \alpha_0 = 0$ , then for all  $m \in N_0$ , we have

$$\begin{cases} x_{4tm+2j+1} &= x_{2j-4t+1}, \\ x_{4tm+2j} &= x_{2j-4t}, \\ y_{4tm+2j+1} &= y_{2j-4t+1}, \\ y_{4tm+2j} &= y_{2j-4t}. \end{cases}$$

#### REFERENCES

- [1] Belhannache, F., Touafek, N. and Abo-Zeid, R., On a higher order rational difference equations, *Journal of Applied Mathematics and Informatics*, 34(5-6), (2016), 369–382.
- [2] Dekkar, I., Touafek, N. and Yazlik, Y., Global stability of a third-order nonlinear system of difference equations with period two coefficients, *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales - Serie A: Matemáticas*, 111, (2017), 325–347.
- [3] Elabbasy, E. M., El-Metwally, H. and Elsayed, E. M., Qualitative behavior of higher order difference equation, *Soochow J. Math.*, 33, (2007), 861–873.
- [4] Cinar, C., Toufik, M. and Yalcinkaya, I., On the difference equation of higher order, *Utilitas Mathematica*, 92, (2013), 161–166.
- [5] Cinar, C., On the positive solutions of the difference equation  $x_{n+1} = \frac{x_{n-1}}{1+x_n x_{n-1}}$ , *Applied Mathematics and Computation*, 150(1), (2004), 21–24.
- [6] Elaydi, S., An Introduction to Difference Equations, Springer, New York, 1996.
- [7] El-Dessoky, M. M., The form of solutions and periodicity for some systems of third-order rational difference equations, *Mathematical Methods in the Applied Sciences*, 39(5), (2016), 1076–1092.
- [8] Elsayed, E. M. and Ahmed, A. M., Dynamics of a three dimensional system of rational difference equations, *Mathematical Methods in the Applied Sciences*, 39(5), (2016), 1026–1038.
- [9] Elsayed, E. M., Alotaibi, A. and Almaylabi, H. A., On a solutions of fourth order rational systems of difference equations, *Journal of Computational Analysis and Applications*, 22(7), (2017), 1298-1308.
- [10] Elsayed, E. M., On the solutions and periodic nature of some systems of difference equations, *International Journal of Biomathematics*, 7(6), (2014), 1–26.
- [11] Gelisken, A. and Kara, M., Some general systems of rational difference equations, *Journal of Difference Equations*, 396757,(2015), 1–7.
- [12] Grove, E. A. and Ladas, G., Periodicities in nonlinear difference equations, *Advances in Discrete Mathematics and Applications*, vol. 4. Chapman & Hall/CRC, London, 2005.
- [13] Haddad, N., Touafek, N. and Rabago, J. F. T., Well-defined solutions of a system of difference equations, *Journal of Appl. Math. and Comput.*, DOI: 10.1007/s12190-017-1081-8, (2017), 1–20.
- [14] Haddad, N., Touafek, N. and Rabago, J.F.T., Solution form of a higher-order system of difference equations and dynamical behavior of its special case, *Mathematical Methods in the Applied Science*, 40(10), (2017), 3599–3607.

- [15] Halim, Y., Touafek, N. and Yazlik, Y., Dynamic behavior of a second-order nonlinear rational difference equation, *Turk. J. Math*, 39(6), (2015), 1004–1018.
- [16] Ibrahim, T.F. and Touafek, N., On a third order rational difference equation with variable coefficients, *Dynamics of Continuous Discrete and Impulsive Systems Series B*, 20, (2013), 251–264.
- [17] Kulenovic, M. R. S. and Ladas, G., Dynamics of Second Order Rational Difference Equations with Open Problems and Conjectures. Chapman & Hall, CRC Press, London, 2001.
- [18] Psarros, N., Papaschinopoulos, G. and Schinas, C.J., Semistability of two systems of difference equations using centre manifold theory, *Mathematical Methods in the Applied Sciences*, 39(18), (2016), 5216–5222.
- [19] Rabago, J.F.T. and Bacani, J.B., On a nonlinear difference equations, *Dynamics of Continuous, Discrete and Impulsive Systems Series A: Mathematical Analysis*, 24, (2017), 375–394.
- [20] Stević, S., More on a rational recurrence relation, *Appl. Math, E-Notes* 4, (2004), 80–85.
- [21] Stević, S., On a system of difference equations with period two coefficients, *Applied Mathematics and Computation*, 218, (2011), 4317–4324.
- [22] Stević, S., On the difference equation  $x_n = \frac{x_{n-2}}{b_n + c_n x_n x_{n-2}}$ , *Appl. Math. Comput.*, 218, (2011), 4507–4513.
- [23] Stević, S., On a solvable rational system of difference equations, *Appl. Math. Comput.* 219, (2012), 2896–2908.
- [24] Stević, S., Representation of solutions of bilinear difference equations in terms of generalized Fibonacci sequences, *Electronic Journal of Qualitative Theory of Differential Equations*, (67), (2014), 1–15.
- [25] Stević, S., Alghamdi, M. A., Shahzad, N. and Maturi, D. A., On a class of solvable difference equations, *Abstract and Applied Analysis*, (2013), Article ID: 157-943, 7 pages.
- [26] Stević, S., On a solvable system of difference equations of  $k$ th order, *Appl. Math. Comput.*, 219, (2013), 7765–7771.
- [27] Stević, S., Diblik, J., Iricanin, B. and Smarda, Z., On a solvable system of rational difference equations, *Journal of Difference Equations and Applications*, 20(5-6), (2014), 811–825.
- [28] Taskara, N., Tollu, D. T. and Yazlik, Y., Solutions of rational difference system of order three in terms of Padovan numbers, *Journal of Advanced Research in Applied Mathematics*, 7(3), (2015), 18–29.
- [29] Tollu, D. T., Yazlik, Y. and Taskara, N., On fourteen solvable systems of difference equations, *Applied Mathematics and Computation*, 233, (2014), 310–319.
- [30] Tollu, D. T., Yazlik, Y. and Taskara, N., Behavior of positive solutions of a difference equation, *Journal of Applied Mathematics and Informatics*, 35, (2017), 217–230.
- [31] Tollu, D.T., Yazlik, Y. and Taskara, N., On the solutions of two special types of Riccati difference equation via Fibonacci numbers, *Advances in Difference Equations*, 2013:174, (2013), 1-7.
- [32] Tollu D. T., Yazlik Y. and Taskara N., On a solvable nonlinear difference equation of higher-order, *Turkish Journal of Mathematics*, 42(4), (2018), 1765-1778.
- [33] Tollu, D. T. and Yalcinkaya, I., Global behavior of a three-dimensional system of difference equations of order three, *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.*, 68(1), (2019), 1–16.
- [34] Touafek, N., On a second order rational difference equation, *Hacet. J. Math. Stat.*, 41, (2012), 867–874.
- [35] Yalcinkaya, I., Hamza, A. E. and Cinar, C., Global behavior of a recursive sequence, *Selçuk Journal of Applied Mathematics*, 14(1), (2013), 3–10.
- [36] Yalcinkaya, I. and Tollu, D. T., Global behavior of a second order system of difference equations, *Advanced Studies in Contemporary Mathematics*, 26(4), (2016), 653–667.
- [37] Yazlik, Y., On the solutions and behavior of rational difference equations, *J. Comput. Anal. Appl.*, 17(3), (2014), 584–594.

- [38] Yazlik, Y., Elsayed, E. M. and Taskara, N., On the behaviour of the solutions the solutions of difference equation system, *J. Comput. Anal. Appl.*, 16(5), (2014), 932–941.
- [39] Yazlik, Y., Tollu, D. T. and Taskara, N., On the behaviour of solutions for some systems of difference equations, *J. Comput. Anal. Appl.*, 18(1), (2015), 166–178.
- [40] Yazlik, Y., Tollu, D. T. and Taskara, N., On the solutions of a three-dimensional system of difference equations, *Kuwait Journal of Science*, 43(1), (2015), 95–111.

*Current address:* Yasin Yazlik: Nevsehir Hacı Bektaş Veli University, Faculty of Science and Art, Department of Mathematics, Nevsehir, Turkey

*E-mail address:* [yyazlik@nevsehir.edu.tr](mailto:yyazlik@nevsehir.edu.tr)

ORCID Address: <http://orcid.org/0000-0001-6369-540X>

*Current address:* Merve KARA: Aksaray University, Ortaköy Vocational School, Aksaray, Turkey

*E-mail address:* [mervekara@aksaray.edu.tr](mailto:mervekara@aksaray.edu.tr)

ORCID Address: <http://orcid.org/0000-0001-8081-0254>